



## Yalamanchili Surgery Packet

Patient: \_\_\_\_\_

DOB: \_\_\_\_\_

Procedure: \_\_\_\_\_

Diagnosis: \_\_\_\_\_

Surgery Date: \_\_\_\_\_

Pre-Operative Appointment: \_\_\_\_\_

Pre-Anesthesia Testing (PAT) Appointment: \_\_\_\_\_

Post-Operative Appointment: \_\_\_\_\_

**Medical City Frisco**  
5500 Frisco Square Blvd  
Frisco, TX 75034  
214-618-0500

**Allen Surgery Center**  
1125 Raintree Circle, Suite 200  
Allen, TX 75013  
469-898-8400

**Live Oak Surgery Center**  
7211 Preston Rd, Suite 2100  
Plano, TX 75024  
469-488-1000

---

### Important Phone Numbers

Office: 972-566-5255

Office Fax: 833-774-2575

Surgery Scheduler: 972-566-5255 (ext. 15110)

Medical Assistant: 972-566-5255 (ext. 15117)

PAT: 214-618-0500

DME: 512-436-9186

---

### Surgery Day Checklist

Medications

DME (sling, crutches, brace, ice machine)

Medical Clearance (PAT, PCP, cardiology)

NPO after midnight on surgery day

Transportation home after surgery

Bring ID, insurance card, and payment

---

## Rehabilitation

Physical therapy begins at 2 weeks after surgery. We usually give you your physical therapy prescription at your initial post-operative visit. Your rehabilitation protocol largely depends on what we end up doing during surgery. We give you our list for recommended facilities, but you can take your prescription anywhere you want.

---

## Wound Care

Arthroscopy Incisions: Change dressings to regular bandages after 24 hours. Keep incisions covered, dry when showering for 2 weeks.

Open Incisions: Keep dressings on for 2 weeks. No need to change anything. Keep incisions covered, dry when showering for 2 weeks.

---

## Return Precautions

### Neurovascular Concerns ⚠️

- Regional nerve block should wear off by 48 hours.
- Watch for significant numbness, tingling, or weakness.
- Watch for abnormal coolness, paleness, or pulselessness.
- Go to ER immediately for any concerns.

### Compartment Syndrome ⚠️

- Watch for severe pain not responding to pain medications.
- Watch for pain with gentle stretching fingers, toes.
- Watch for significant swelling, tightness, or blistering.
- Watch for significant numbness, tingling, or weakness.
- Go to ER immediately for any concerns.

### Blood Clot (DVT) ⚠️

- Watch for increased posterior calf, thigh pain.
- Watch for abnormal unilateral leg swelling.
- Watch for difficulty breathing at rest. Sometimes it occurs even with light activity.
- Go to ER immediately for any concerns.

### Pulmonary Embolus (PE) ⚠️

- Watch for sudden difficulty breathing.

- Watch for chest pain with deep breathing.
- Watch for rapid heart rate.
- Watch for lightheadedness, fainting, or vision problems.
- Go to ER immediately for any concerns.

#### Wound Concerns

- Watch for dressings become wet.
- Watch for persistent bleeding that soaks through dressings.
- Watch for wound dehiscence.
- Call office for any questions.
- Go to ER immediately for any concerns.

#### Infection

- Watch for fever greater than 101°F not responding to Tylenol.
- Watch for worsening redness, swelling, or pus around incision.
- Call office for any questions.
- Go to ER immediately for any concerns.

#### Medication Allergy

- Most common allergies include reactions to skin glue, dressings, or cleaning solution used during surgery.
- Watch for rash, itching, or hives not responding to Benadryl.
- Watch for severe reactions, including difficulty breathing, face swelling, or difficulty swallowing.
- Call office for any questions.
- Go to ER immediately for any concerns.

#### General Concerns

- Watch for worsening pain not responding to medications.
- Watch for fever greater than 101°F not responding to Tylenol.
- Watch for difficulty breathing, swallowing, or talking.
- Please call office for any symptoms that appear worrisome.

## About Dr. Raj Yalamanchili

Dr. Raj Yalamanchili is a double board-certified orthopaedic surgeon with advanced fellowship training in sports medicine, shoulder replacement, and minimally invasive, arthroscopic treatment of the shoulder, hip, and knee. He also offers various conservative treatment options, including cortisone, hyaluronic acid, and PRP injections under ultrasound guidance.

Dr. Yalamanchili is a native of Plano, TX, where he graduated from Plano Senior High School in 2007. He earned his bachelor's degree from the University of Southern California (USC) in 2011, before graduating with Distinction in Research from the Keck School of Medicine of USC in 2015.

He then went on to complete his orthopaedic residency at Cedars-Sinai Medical Center in Los Angeles, CA in 2020. Dr. Yalamanchili elected to further his training by completing a fellowship in sports medicine at the Kerlan-Jobe Institute in Los Angeles, CA. During his fellowship, Dr. Yalamanchili served as assistant team physician for the Los Angeles Rams (NFL), Los Angeles Dodgers (MLB), Los Angeles Sparks (WNBA), and Los Angeles Galaxy (MLS).

After fellowship, Dr. Yalamanchili took a position as Assistant Professor at Loma Linda University in Loma Linda, CA. He also served as Medical Director at SAC Health System, a nonprofit community healthcare clinic in San Bernardino, CA. During his time at Loma Linda University, Dr. Yalamanchili was the head team physician for the Rancho Cucamonga Quakes baseball team, a Single-A affiliate of the Los Angeles Dodgers.

Dr. Yalamanchili is a fellow of the American Academy of Orthopaedic Surgeons (FAAOS), and he also holds memberships with the American Orthopaedic Society of Sports Medicine (AOSSM) and the Arthroscopy Association of North America (AANA). Dr. Yalamanchili believes in empowering patients through empathy and compassion, while embracing innovative techniques and technology throughout the healing process to achieve the best possible outcome.



## Medications

### Shoulder

Voltaren 100 mg XR PO daily x 14 days, #14 pills  
Lyrica 75 mg PO daily x 14 days, #14 pills  
Percocet 10/325 mg PO q8h as needed for pain, #20 pills

---

### Elbow

Voltaren 100 mg XR PO daily x 14 days, #14 pills  
Lyrica 75 mg PO daily x 14 days, #14 pills  
Percocet 10/325 mg PO q8h as needed for pain, #20 pills

---

### Hip

Ecotrin 325 mg PO daily x 42 days, #42 pills  
Losartan 25 mg PO daily x 6 weeks, #42 pills  
Voltaren 100 mg XR PO daily x 14 days, #14 pills  
Lyrica 75 mg PO daily x 14 days, #14 pills  
Percocet 10/325 mg PO q8h as needed for pain, #20 pills

---

### Knee

Ecotrin 325 mg PO daily x 14 days, #14 pills  
Voltaren 100 mg XR PO daily x 14 days, #14 pills  
Lyrica 75 mg PO daily x 14 days, #14 pills  
Percocet 10/325 mg PO q8h as needed for pain, #20 pills

---

### Ankle

Ecotrin 325 mg PO daily x 14 days, #14 pills  
Voltaren 100 mg XR PO daily x 14 days, #14 pills  
Lyrica 75 mg PO daily x 14 days, #14 pills  
Percocet 10/325 mg PO q8h as needed for pain, #20 pills

---

## Stop Medications Before Surgery\*

Medication Class	Examples	Stop Before Surgery	Notes
Warfarin	Coumadin	5 days	Goal INR <1.5
DOAC	Apixaban, Rivaroxaban, Dabigatran	3–5 days	Longer if renal impaired
Aspirin	ASA, Ecotrin	5–7 days	Discuss with cardiology for stents

Medication Class	Examples	Stop Before Surgery	Notes
Clopidogrel	Plavix	5–7 days	Discuss with cardiology for stents
NSAID	Ibuprofen, Naproxen	3–7 days	Increased bleeding risk
Biologics (TNF Inhibitors)	Humira, Enbrel, Remicade	1 dosing cycle before	Increased infection risk
JAK Inhibitors	Tofacitinib, Upadacitinib	3–7 days	Discuss with rheumatology
Methotrexate	MTX	Continue	Consider holding with renal issues
Chronic Steroids	Prednisone	Continue	Might need stress dosing
GLP-1 Agonists	Semaglutide (Ozempic), Tirzepatide	7 days	Aspiration risk
Metformin	Glucophage	Hold on morning	Resume if renal function stable
Insulin	Various	Adjust dose	Discuss with PCP
ACEI / ARB	Lisinopril, Losartan	Hold on morning	To prevent hypotension
Beta Blockers	Metoprolol	Continue	To avoid withdrawal
Supplements / Herbals	Fish oil, Vitamin E, ginkgo	7–14 days	Increased bleeding risk

\*Please check with your prescribing physician before stopping any medications.

### Resume Medications After Surgery

Medication Class	Resume
Anticoagulants	24 hours after surgery
Biologics	2 weeks after surgery
GLP-1	24 hours after surgery
ACEI/ARB	24 hours after surgery
Metformin	24 hours after surgery